

REFERENCES

- Bagyaraj, M., Gurugnanam, B., & Nagar, A. (2011). Significance of morphometry studies, soil characteristics, erosion phenomena and landform processes using remote sensing and GIS for Kodaikanal Hills, a global biodiversity hotspot in Western Ghats, Dindigu District, Tamil Nadu, South India. *Research Journal of Environmental and Earth Sciences*, 3(3), 221-233.
- Bapalu, G. V., & Sinha, R. (2005). GIS in flood hazard mapping: A case study of Kosi River Basin, India. *GIS Development Weekly*, pp. 1-3.
- Bhatt, S., & Ahmed, S. A. (2014). Morphometric analysis to determine floods in the Upper Krishna basin using Cartosat DEM. *Geocarto International*, 29(8), 878-894.
- El Maghraby, M., Masoud, M., & Niyazi, B. (2014). Assessment of Surface Runoff in Arid, Data Scarce Regions; An Approach Applied in Wadi Al Hamd, Al Madinal al Munawarah, Saudi Arabia. *Life Science Journal*, 11(4).
- El-Behiry, M. G., Shedid, A., & Abu-Khadra, A. (2006). Integrated GIS and remote sensing for runoff hazard analysis in AIn Sukhna industrail area, Egypt. *Earth Sciences*, 19-41.
- Elmoustafa, A. M., & Mona, M. M. (2013). lash Flood Risk Assessment Using Morphological Parameters in Sinai Peninsula. *Open Journal of Modern Hydrology*.
- Elmoustafa, M. A. (2012). Weighted normalized risk factor for floods risk assessment. *Ain Shams Engineering Journal*, 327–332.
- FloodList. (2017). Asia. Retrieved September 09, 2017, from Reporting floods and flooding news since 2008: <http://floodlist.com/asia/pakistan-gilgit-baltistan-august-2017>
- Javed, A., Ahmad, R., & Khanday, M. Y. (2009). Prioritization of Sub-watersheds based on Morphometric and Land Use Analysis using Remote Sensing and GIS Techniques. *Journal of Idian*, 261-274.
- Omran, A., Schroder, D., EL, R. A., & Geriesh, M. (2011). Flood Hazard Assessment in Wadi Dahab, Egypt Based on Basin Morphometry Using GIS Techniques. *GI_Forum Program Committee*.
- Patton, C. P., & Baker, R. V. (1976). Morphometry and Floods in Small Drainage Basins Subject to Diverse Hydrogeomorphic Controls. *Water Resources Research*, 12.
- Patton, P. C. (1988). Drainage basin morphometry and floods. In *Flood Geomorphology* (pp. 51-64). New York: John Wiley & Sons
- Pisal, P. A., Yadav, A. S., & Chavan, A. (2013). Morphometric Analysis Of Bhogavati River Basin, Kolhapur District, Maharashtra, India. *Journal of Mechanical and Civil Engineering* (, 01-08.
- Strahler, A. N. (1957). Quantitative Analysis of Watershed Geomorphology. *Transaction American Geophysical Union*, Vol 38, No 6.
- Subyani, A. M., & Al-Ahmadi, F. S. (2011). RAINFALL-RUNOFF MODELING IN THE AL-MADINAH AREA OF WESTERN SAUDI ARABIA. *Journal of Environmental Hydrology*.
- Sujata, B., Sudhakar, S., & V R, D. (1999). Prioritisation of sub watershed based on morphometric analysis of drainage basin; A remote sensing and GIS approach. *Journal of the Indian Society of Remote Sensing*, Vol. 27 No. 3.
- The Express Tribune Blogs. (2016, Feb 15). Retrieved from The Express Tribune: <http://tribune.com.pk/story/933507/over-a-million-people-affected-by-floods-in-pakistan-ndma/>
- Waikar, M. L., & Nilawar, A. P. (2014). Morphometric Analysis of a Drainage Basin Using Geographical Information System: A Case study. *International Journal of Multidisciplinary and Current Research*, 2.
- World Wild Life. (Dec 2009). Land cover mapping of Central Karakoram National Park. *GIS Laboratories*.
- Youssef, A. M., Pradhan, B., & Hassan, A. M. (2011). Flash flood risk estimation along the St. Katherine road, southern Sinai, Egypt using GIS based morphometry and satellite imagery. *Environmental Earth Sciences*, 62(3), 611-623.